Barriers to cervical screening in women who have experienced sexual abuse: an exploratory study

Web-only data (Table S3) – Measures of association with ever attended for cervical screening

	Ever atte	nded for smea	r
	No	Yes	Fisher's exact (p)
Barriers			
Self-worth			
Not coded	17	74	1.00
Coded	4	22	
Power disparity			
Not coded	14	69	0.61
Coded	7	27	
Trust, safety, disclosure			
Not coded	17	76	1.00
Coded	4	20	
Physical pain			
Not coded	20	71	0.041
Coded	1	25	
Sexual victimisation			
Not coded	13	59	1.00
Coded	8	37	
Mechanics of examination			
Not coded	18	88	0.41
Coded	3	8	

	Ever attended for smear		
	No	Yes	Fisher's exact (p)
Fear and anxiety			
Not coded	9	71	0.009
Coded	12	25	
Communication barrier			
Not coded	19	78	0.522
Coded	2	18	
mpact on coping strategies			
Not coded	12	58	0.81
Coded	58	38	
Intention to attend			
Not coded	12	67	0.31
Coded	9	29	
mprovement of cervical screening			
Disclosure			
Not coded	18	77	0.76
Coded	3	19	
Safety, trust and sharing control			
Not coded	13	52	0.63
Coded	8	44	
Communication			
Not coded	11	44	0.64
Coded	10	52	
Position, procedure, environment			
Not coded	18	76	0.76
Coded	3	20	

	Ever attended for smear		
	No	Yes	Fisher's exact (p)
Time and space			
Not coded	15	65	0.80
Coded	6	31	
Relationship status			
Single	9	32	0.59
In a relationship	4	13	
Married/living with partner	9	47	
Separated/divorced	0	10	
Widowed	0	1	
Do not wish to answer	0	2	
Age left education			
≤15 years	2	13	0.59
16 years	2	15	
17 years	0	10	
18 years	5	15	
≥19 years	13	51	
Do not wish to answer	0	1	
Physical abuse			
Yes	8	55	0.24
No	14	50	
Emotional abuse			
Yes	6	35	0.80
No	16	70	

	Ever att	Ever attended for smear			
	No	Yes	Fisher's exact (p)		
Neglect					
Yes	15	79	0.59		
No	7	26			
Ritual abuse					
Yes	19	100	0.14		
No	3	5			
Spiritual abuse					
Yes	21	95	0.69		
No	1	10			
Other abuse					
Yes	20	101	0.28		
No	2	4			